DEPARTMENT OF HEALTH

STI Quarterly Surveillance Report

JAN. 1 – JUNE 30, 2026

Overview

Table 1. Number of Cases and Rates (per 100,000 persons) of Chlamydia, Gonorrhea, Syphilis, and Chancroid – Minnesota, 2022 – Q1-Q2 2026

Disease	2022 Cases	2022 Rate	2023 Cases	2023 Rate	2024 Cases	2024 Rate	2025 Cases	2025 Rate	Q1-Q2 2026 Cases	Q1-Q2 2026 Rate
Chlamydia	22,079	386.7	21,767	380.7	19,703	344.6	18,179	318.0	8,630	150.9
Gonorrhea	8,161	153.9	7,717	135.0	7,131	124.7	5,960	104.2	2,833	49.6
All Stages of Syphilis	1831	34.5	1748	30.6	1478	25.9	1384	24.2	524	9.2
Primary/Secondary Syphilis	676	12.7	502	8.8	440	7.7	373	6.5	144	2.5
Early NP/NS	535	10.1	387	6.8	755	13.2	299	5.2	80	1.4
Late Unknown Duration[I]	601	11.3	830	14.5	694	12.1	682	11.9	286	5.0
Congenital Syphilis	20	31.7	26	42.7	29	47.5	30	47.5	14	23.0
Chancroid	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

Note: Data exclude cases diagnosed in federal or private correctional facilities.
U.S. Census Intercensal and U.S. 2020/2022 data is used to calculate rates.
[I]Includes unstaged neurosyphilis, latent syphilis of unknown duration, and late syphilis with clinical manifestations.
[II]Congenital syphilis rate per 100,000 live births.

Chlamydia

Note: Data exclude cases diagnosed in federal or private correctional facilities.

U.S. Census Intercensal and U.S. 2020/2022 data is used to calculate rates.

Table 2.A. Number of Cases and Rates (per 100,000 persons) of Chlamydia by Residence and Gender – Minnesota, Q1-Q2 2026

Residence[I]	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
Minneapolis	752	25%	1,054	19%	1,809	21%	426
St. Paul	315	10%	679	12%	998	12%	329
Suburban[II]	1,140	37%	2,047	37%	3,189	37%	132
Greater Minnesota	830	27%	1,743	31%	2,578	30%	100

Residence[I]	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
Unknown[III]	24	1%	32	1%	56	1%	x
Total[IV]	3,061	100%	5,555	100%	8,630	100%	*151*

[I]Residence missing for 56 cases of chlamydia.
[II]Suburban is defined as the seven-county metropolitan area (Anoka, Carver, Dakota, Hennepin, Ramsey, Scott, and Washington Counties, excluding the cities of Minneapolis and St. Paul).
[III]No comparable population data available to calculate rates.
[IV]Gender missing/other/unknown for 14 cases of chlamydia.

Table 2.B. Number of Cases and Rates of Chlamydia by Age and Gender – Minnesota, Q1-Q2 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
< 15 yrs	11	0%	44	1%	55	1%	8
15-19 yrs	591	19%	1,744	31%	2,338	27%	625
20-24 yrs	994	32%	1,970	35%	2,969	34%	800
25-29 yrs	535	17%	875	16%	1,412	16%	390
30-34 yrs	332	11%	423	8%	757	9%	200
35-39 yrs	239	8%	233	4%	472	5%	120
40-44 yrs	150	5%	134	2%	285	3%	75
45-49 yrs	91	3%	76	1%	167	2%	52
50-54 yrs	41	1%	26	0%	67	1%	20
55+ yrs	77	3%	30	1%	108	1%	6
Total[I]	3,061	100%	5,555	100%	8,630	100%	*151*

[I]Gender missing/other/unknown for 14 cases of chlamydia.

Table 2.C. Number of Cases and Rates of Chlamydia by Race/Ethnicity and Gender – Minnesota, Q1-Q2 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
White, Non-Hispanic	845	28%	1,827	33%	2,673	31%	60
Black, Non-Hispanic	1,088	36%	1,463	26%	2,554	30%	604
American Indian	61	2%	216	4%	277	3%	462
Asian/PI	96	3%	235	4%	334	4%	107
Other[I,II]	158	5%	316	6%	474	5%	x

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
Unknown[II]	483	16%	787	14%	1,273	15%	x
Hispanic[IV]	330	11%	711	13%	1,045	12%	306
Total[IV]	3,061	100%	5,555	100%	8,630	100%	*151*

[I]Includes persons reported with more than one race.
[II]No comparable population data available to calculate rates.
[III]Persons of Hispanic origin may be of any race.
[IV]Gender missing/other/unknown for 14 cases of chlamydia.

Gonorrhea

Note: Data exclude cases diagnosed in federal or private correctional facilities.

U.S. Census Intercensal and U.S. 2020/2022 data is used to calculate rates.

Table 3.A. Number of Cases and Rates (per 100,000 persons) of Gonorrhea by Residence and Gender – Minnesota, Q1-Q2 2026

Residence[I]	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
Minneapolis	735	41%	357	35%	1,102	39%	259
St. Paul	278	15%	156	15%	436	15%	144
Suburban[II]	582	32%	341	34%	929	33%	38
Greater Minnesota	190	11%	155	15%	345	12%	13
Unknown[III]	16	1%	5	0%	21	1%	x
Total[IV]	1,801	100%	1,014	100%	2,833	100%	50

[I]Residence missing for 21 cases of gonorrhea.
[II]Suburban is defined as the seven-county metropolitan area (Anoka, Carver, Dakota, Hennepin, Ramsey, Scott, and Washington Counties, excluding the cities of Minneapolis and St. Paul).
[III]No comparable population data available to calculate rates.
[IV]Gender missing/other/unknown for 18 cases of gonorrhea.

Table 3.B. Number of Cases and Rates of Gonorrhea by Age and Gender – Minnesota, Q1-Q2 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
< 15 yrs	5	0%	11	1%	16	1%	2
15-19 yrs	132	7%	222	22%	356	13%	95
20-24 yrs	326	18%	260	26%	592	21%	160
25-29 yrs	308	17%	158	16%	468	17%	129
30-34 yrs	337	19%	131	13%	471	17%	124
35-39 yrs	260	14%	106	10%	369	13%	94
40-44 yrs	159	9%	67	7%	227	8%	60
45-49 yrs	101	6%	33	3%	135	5%	42
50-54 yrs	65	4%	17	2%	82	3%	24
55+ yrs	108	6%	9	1%	117	4%	7
Total[I]	1,801	100%	1,014	100%	2,833	100%	50

[I]Gender missing/other/unknown for 18 cases of gonorrhea.

Table 3.C. Number of Cases and Rates of Gonorrhea by Race/Ethnicity and Gender – Minnesota, Q1-Q2 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
White, Non-Hispanic	545	30%	288	28%	841	30%	19
Black, Non-Hispanic	697	39%	367	36%	1,068	38%	252
American Indian	25	1%	71	7%	96	3%	160
Asian/PI	57	3%	20	2%	79	3%	25
Other[I,II]	113	6%	56	6%	170	6%	x
Unknown[II]	215	12%	137	14%	354	12%	x
Hispanic[III]	149	8%	75	7%	225	8%	66
Total[IV,V]	1,801	100%	1,014	100%	2,833	100%	50

[I]Includes persons reported with more than one race.
[II]No comparable population data available to calculate rates.
[III]Persons of Hispanic origin may be of any race.
[IV]Gender missing/other/unknown for 18 cases of gonorrhea.

Primary/Secondary Syphilis

Note: Data exclude cases diagnosed in federal or private correctional facilities.

U.S. Census Intercensal and U.S. 2022 data is used to calculate rates.

Table 4.A. Number of Cases and Rates (per 100,000 persons) of Primary/Secondary Syphilis by Residence and Gender – Minnesota, Q1-Q2 2026

Residence	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
Minneapolis	36	34%	12	32%	49	34%	11.5
St. Paul	17	16%	2	5%	19	13%	6.3
Suburban[I]	27	26%	14	37%	41	28%	1.7
Greater Minnesota	25	24%	10	26%	35	24%	1.4
Total[II]	105	100%	38	100%	144	100%	2.5

[I]Suburban is defined as the seven-county metropolitan area (Anoka, Carver, Dakota, Hennepin, Ramsey, Scott, and Washington Counties, excluding the cities of Minneapolis and St. Paul).
[II]Gender missing/other/unknown for 1 case of syphilis.

Table 4.B. Number of Cases and Rates of Primary/Secondary Syphilis by Age and Gender – Minnesota, Q1-Q2 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
< 15 yrs	0	0%	0	%	0	0%	0.0
15-19 yrs	2	2%	1	3%	3	2%	0.8
20-24 yrs	8	8%	5	13%	13	9%	3.5
25-29 yrs	16	15%	11	29%	27	19%	7.5
30-34 yrs	17	16%	6	16%	24	17%	6.3
35-39 yrs	15	14%	4	11%	19	13%	4.8
40-44 yrs	8	8%	5	13%	13	9%	3.4
45-49 yrs	14	13%	3	8%	17	12%	5.3
50-54 yrs	6	6%	2	5%	8	6%	2.4
55+ yrs	19	18%	1	3%	20	14%	1.2
Total[I]	105	100%	38	100%	144	100%	2.5

[I]Gender missing/other/unknown for 1 case of syphilis.

Table 4.C. Number of Cases and Rates of Primary/Secondary Syphilis by Race/Ethnicity and Gender – Minnesota, Q1-Q2 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
White, Non-Hispanic	24	23%	11	29%	35	24%	0.8
Black, Non-Hispanic	44	42%	14	37%	59	41%	13.9
American Indian	4	4%	4	11%	8	6%	13.4
Asian/PI	6	6%	2	5%	8	6%	2.6
Other[I,II]	6	6%	3	8%	9	6%	x
Unknown[II]	0	0%	1	3%	1	1%	x
Hispanic[III]	21	20%	3	8%	24	17%	7.0
Total[IV]	105	100%	38	100%	144	100%	2.5

[I]Includes persons reported with more than one race.
[II]No comparable population data available to calculate rates.
[III]Persons of Hispanic origin may be of any race.
[IV]Gender missing/other/unknown for 1 case of syphilis.

County STI Totals

Note: Data exclude cases diagnosed in federal or private correctional facilities.

County data missing for 56 chlamydia cases and 22 gonorrhea cases.

U.S. Census Intercensal and U.S. 2022 data is used to calculate rates.

Table 5. Number of Cases and Rates[I] (per 100,000 persons) of Chlamydia and Gonorrhea by County of Residence – Minnesota, Q1-Q2 2026

County	Cases (Chlamydia)	Rate (Chlamydia)	Cases (Gonorrhea)	Rate (Gonorrhea)
Aitkin	6	37	0	-
Anoka	510	138	127	34
Becker	37	105	2	-
Beltrami	79	169	7	15
Benton	46	111	7	17
Big Stone	5	97	2	-
Blue Earth	119	171	11	16
Brown	14	54	2	-
Carlton	34	93	4	-

County	Cases (Chlamydia)	Rate (Chlamydia)	Cases (Gonorrhea)	Rate (Gonorrhea)
Carver	76	69	13	12
Cass	29	93	2	-
Chippewa	13	106	0	-
Chisago	47	81	4	-
Clay	120	182	24	36
Clearwater	6	69	0	-
Cook	3	-	1	-
Cottonwood	16	141	0	-
Crow Wing	50	74	11	16
Dakota	589	133	147	33
Dodge	21	100	2	-
Douglas	21	53	3	-
Faribault	8	57	2	-
Fillmore	13	61	2	-
Freeborn	42	137	1	-
Goodhue	37	77	10	21
Grant	3	-	0	-
Hennepin	3,040	241	1544	123
Houston	9	48	2	-
Hubbard	18	82	0	-
Isanti	37	87	6	14
Itasca	41	91	1	-
Jackson	7	71	0	-
Kanabec	21	128	0	-
Kandiyohi	51	116	5	11
Kittson	5	123	0	-
Koochiching	4	-	1	-
Lac qui Parle	5	75	0	-
Lake	6	55	0	-
Lake of the Woods	2	-	0	-
Le Sueur	20	69	5	17
Lincoln	2	-	0	-

County	Cases (Chlamydia)	Rate (Chlamydia)	Cases (Gonorrhea)	Rate (Gonorrhea)
Lyon	24	95	0	-
McLeod	27	74	2	-
Mahnomen	11	206	0	-
Marshall	4	-	0	-
Martin	10	51	1	-
Meeker	10	43	2	-
Mille Lacs	19	70	2	-
Morrison	18	53	6	18
Mower	75	187	8	20
Murray	6	74	0	-
Nicollet	21	61	6	17
Nobles	57	260	3	-
Norman	4	-	2	-
Olmsted	220	134	26	16
Otter Tail	36	59	5	8
Pennington	10	72	2	-
Pine	22	75	1	-
Pipestone	8	86	0	-
Polk	32	104	5	16
Pope	5	44	0	-
Ramsey	1,275	238	517	96
Red Lake	2	-	0	-
Redwood	9	59	0	-
Renville	8	55	0	-
Rice	52	77	14	21
Rock	6	63	2	-
Roseau	6	39	3	-
St. Louis	196	98	51	26
Scott	198	128	53	34
Sherburne	105	104	18	18
Sibley	10	67	0	-
Stearns	223	139	31	19

County	Cases (Chlamydia)	Rate (Chlamydia)	Cases (Gonorrhea)	Rate (Gonorrhea)
Steele	47	126	5	13
Stevens	10	104	3	-
Swift	12	123	1	-
Todd	24	94	2	-
Traverse	2	-	0	-
Wabasha	12	55	0	-
Wadena	9	63	2	-
Waseca	18	95	2	-
Washington	299	108	62	22
Watonwan	10	90	2	-
Wilkin	4	-	1	-
Winona	55	111	2	-
Wright	143	97	21	14
Yellow Medicine	8	84	0	-

[l]Rates not calculated for counties with fewer than five cases.

Note: County syphilis data is available at Newly Diagnosed Syphilis Cases by County (https://www.health.state.mn.us/diseases/stds/stats/county.html).

Minnesota Department of Health
651-201-5414
www.health.state.mn.us/std

10/28/2026

To obtain this information in a different format, call: 651-201-5414.