DEPARTMENT OF HEALTH

STI Quarterly Surveillance Report

JAN. 1 – SEPT. 30, 2026

Overview

Table 1. Number of Cases and Rates (per 100,000 persons) of Chlamydia, Gonorrhea, Syphilis, and Chancroid – Minnesota, 2022 – Q1-Q3 2026

Disease	2022 Cases	2022 Rate	2023 Cases	2023 Rate	2024 Cases	2024 Rate	2025 Cases	2025 Rate	Q1-Q3 2026 Cases	Q1-Q3 2026 Rate
Chlamydia	22,079	386.7	21,767	380.7	19,703	344.6	18,179	318.0	13,092	229.0
Gonorrhea	8,161	153.9	7,717	135.0	7,131	124.7	5,960	104.2	4,487	78.5
All Stages of Syphilis	1831	34.5	1748	30.6	1478	25.9	1384	24.2	750	13.1
Primary/Secondary Syphilis	676	12.7	502	8.8	440	7.7	373	6.5	203	3.6
Early NP/NS	535	10.1	387	6.8	755	13.2	299	5.2	123	2.2
Late Unknown Duration[I]	601	11.3	830	14.5	694	12.1	682	11.9	403	7.0
Congenital Syphilis[II]	20	31.7	26	42.7	29	47.5	30	47.5	21	34.5
Chancroid	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

Note: Data exclude cases diagnosed in federal or private correctional facilities.
U.S. Census Intercensal and U.S. 2020/2022 data is used to calculate rates.
[I]Includes unstaged neurosyphilis, latent syphilis of unknown duration, and late syphilis with clinical manifestations.
[II]Congenital syphilis rate per 100,000 live births.

Chlamydia

Note: Data exclude cases diagnosed in federal or private correctional facilities.

U.S. Census Intercensal and U.S. 2020/2022 data is used to calculate rates.

Table 2.A. Number of Cases and Rates (per 100,000 persons) of Chlamydia by Residence and Gender – Minnesota, Q1-Q3 2026

Residence[I]	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
Minneapolis	1,082	23%	1,512	18%	2,607	20%	613
St. Paul	500	11%	983	12%	1,487	11%	490
Suburban[II]	1,838	39%	3,209	38%	5,057	39%	209
Greater Minnesota	1,221	26%	2,627	31%	3,856	29%	150

Residence[I]	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
Unknown[III]	34	1%	50	1%	85	1%	x
Total[IV]	4,675	100%	8,381	100%	13,092	100%	229

[I]Residence missing for 85 cases of chlamydia.
[II]Suburban is defined as the seven-county metropolitan area (Anoka, Carver, Dakota, Hennepin, Ramsey, Scott, and Washington Counties, excluding the cities of Minneapolis and St. Paul).
[III]No comparable population data available to calculate rates.
[IV]Gender missing/other/unknown for 36 cases of chlamydia.

Table 2.B. Number of Cases and Rates of Chlamydia by Age and Gender – Minnesota, Q1-Q3 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
< 15 yrs	13	0%	67	1%	80	1%	11
15-19 yrs	952	20%	2,595	31%	3,553	27%	949
20-24 yrs	1,475	32%	2,998	36%	4,486	34%	1,209
25-29 yrs	804	17%	1,293	15%	2,104	16%	582
30-34 yrs	513	11%	648	8%	1,168	9%	308
35-39 yrs	354	8%	368	4%	722	6%	183
40-44 yrs	226	5%	214	3%	441	3%	117
45-49 yrs	147	3%	114	1%	262	2%	82
50-54 yrs	64	1%	43	1%	107	1%	31
55+ yrs	127	3%	41	0%	169	1%	10
Total[I]	4,675	100%	8,381	100%	13,092	100%	229

[I]Gender missing/other/unknown for 36 cases of chlamydia.

Table 2.C. Number of Cases and Rates of Chlamydia by Race/Ethnicity and Gender – Minnesota, Q1-Q3 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
White, Non-Hispanic	1,290	28%	2,778	33%	4,076	31%	92
Black, Non-Hispanic	1,668	36%	2,205	26%	3,880	30%	917
American Indian	93	2%	308	4%	402	3%	671
Asian/PI	150	3%	353	4%	507	4%	162
Other[I,II]	244	5%	484	6%	729	6%	x

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
Unknown[II]	712	15%	1,141	14%	1,859	14%	x
Hispanic[IV]	518	11%	1,112	13%	1,639	13%	480
Total[IV]	4,675	100%	8,381	100%	13,092	100%	229

[I]Includes persons reported with more than one race.
[II]No comparable population data available to calculate rates.
[III]Persons of Hispanic origin may be of any race.
[IV]Gender missing/other/unknown for 36 cases of chlamydia.

Gonorrhea

Note: Data exclude cases diagnosed in federal or private correctional facilities.

U.S. Census Intercensal and U.S. 2020/2022 data is used to calculate rates.

Table 3.A. Number of Cases and Rates (per 100,000 persons) of Gonorrhea by Residence and Gender – Minnesota, Q1-Q3 2026

Residence[I]	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
Minneapolis	1,157	41%	559	35%	1,733	39%	408
St. Paul	415	15%	248	15%	666	15%	220
Suburban[II]	948	33%	540	33%	1,498	33%	62
Greater Minnesota	296	10%	265	16%	562	13%	22
Unknown[III]	21	1%	6	0%	28	1%	x
Total[IV]	2,837	100%	1,618	100%	4,487	100%	78

[I]Residence missing for 28 cases of gonorrhea.
[II]Suburban is defined as the seven-county metropolitan area (Anoka, Carver, Dakota, Hennepin, Ramsey, Scott, and Washington Counties, excluding the cities of Minneapolis and St. Paul).
[III]No comparable population data available to calculate rates.
[IV]Gender missing/other/unknown for 32 cases of gonorrhea.

Table 3.B. Number of Cases and Rates of Gonorrhea by Age and Gender – Minnesota, Q1-Q3 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
< 15 yrs	9	0%	14	1%	23	1%	3
15-19 yrs	219	8%	340	21%	562	13%	150
20-24 yrs	492	17%	418	26%	920	21%	248
25-29 yrs	480	17%	265	16%	751	17%	208
30-34 yrs	515	18%	205	13%	727	16%	192
35-39 yrs	421	15%	159	10%	583	13%	148
40-44 yrs	256	9%	114	7%	372	8%	98
45-49 yrs	174	6%	59	4%	234	5%	73
50-54 yrs	97	3%	31	2%	128	3%	38
55+ yrs	174	6%	13	1%	187	4%	11
Total[I]	2,837	100%	1,618	100%	4,487	100%	78

[I]Gender missing/other/unknown for 32 cases of gonorrhea.

Table 3.C. Number of Cases and Rates of Gonorrhea by Race/Ethnicity and Gender – Minnesota, Q1-Q3 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
White, Non-Hispanic	826	29%	480	30%	1,321	29%	30
Black, Non-Hispanic	1,113	39%	567	35%	1,686	38%	398
American Indian	45	2%	122	8%	168	4%	280
Asian/PI	110	4%	46	3%	158	4%	51
Other[I,II]	194	7%	93	6%	288	6%	x
Unknown[II]	330	12%	201	12%	536	12%	x
Hispanic[III]	219	8%	109	7%	330	7%	97
Total[IV,V]	2,837	100%	1,618	100%	4,487	100%	78

[I]Includes persons reported with more than one race.
[II]No comparable population data available to calculate rates.
[III]Persons of Hispanic origin may be of any race.
[IV]Gender missing/other/unknown for 32 cases of gonorrhea.

Primary/Secondary Syphilis

Note: Data exclude cases diagnosed in federal or private correctional facilities.

U.S. Census Intercensal and U.S. 2022 data is used to calculate rates.

Table 4.A. Number of Cases and Rates (per 100,000 persons) of Primary/Secondary Syphilis by Residence and Gender – Minnesota, Q1-Q3 2026

Residence	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
Minneapolis	48	33%	7	13%	67	33%	15.8
St. Paul	22	15%	1	2%	27	13%	8.9
Suburban[I]	42	29%	5	9%	62	31%	2.6
Greater Minnesota	35	24%	6	11%	47	23%	1.8
Total[II]	147	100%	54	100%	203	100%	3.6

[I]Suburban is defined as the seven-county metropolitan area (Anoka, Carver, Dakota, Hennepin, Ramsey, Scott, and Washington Counties, excluding the cities of Minneapolis and St. Paul).
[II]Gender missing/other/unknown for 2 cases of syphilis.

Table 4.B. Number of Cases and Rates of Primary/Secondary Syphilis by Age and Gender – Minnesota, Q1-Q3 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
< 15 yrs	0	0%	0	%	0	0%	0.0
15-19 yrs	2	1%	3	6%	5	2%	1.3
20-24 yrs	10	7%	7	13%	17	8%	4.6
25-29 yrs	20	14%	12	22%	33	16%	9.1
30-34 yrs	23	16%	7	13%	31	15%	8.2
35-39 yrs	26	18%	6	11%	32	16%	8.1
40-44 yrs	16	11%	10	19%	26	13%	6.9
45-49 yrs	19	13%	5	9%	24	12%	7.5
50-54 yrs	8	5%	3	6%	11	5%	3.2
55+ yrs	23	16%	1	2%	24	12%	1.4
Total[I]	147	100%	54	100%	203	100%	3.6

[I]Gender missing/other/unknown for 2 cases of syphilis.

Table 4.C. Number of Cases and Rates of Primary/Secondary Syphilis by Race/Ethnicity and Gender – Minnesota, Q1-Q3 2026

Age	Cases (Males)	% (Males)	Cases (Females)	% (Females)	Cases (Total)	% (Total)	Rate (Total)
White, Non-Hispanic	38	26%	16	30%	55	27%	1.2
Black, Non-Hispanic	60	41%	18	33%	79	39%	18.7
American Indian	4	3%	6	11%	10	5%	16.7
Asian/PI	10	7%	3	6%	13	6%	4.2
Other[I,II]	7	5%	3	6%	10	5%	x
Unknown[II]	0	0%	2	4%	2	1%	x
Hispanic[III]	28	19%	6	11%	34	17%	10.0
Total[IV]	147	100%	54	100%	203	100%	3.6

[I]Includes persons reported with more than one race.
[II]No comparable population data available to calculate rates.
[III]Persons of Hispanic origin may be of any race.
[IV]Gender missing/other/unknown for 2 cases of syphilis.

County STI Totals

Note: Data exclude cases diagnosed in federal or private correctional facilities.

County data missing for 94 chlamydia cases and 42 gonorrhea cases.

U.S. Census Intercensal and U.S. 2022 data is used to calculate rates.

Table 5. Number of Cases and Rates[I] (per 100,000 persons) of Chlamydia and Gonorrhea by County of Residence – Minnesota, Q1-Q3 2026

County	Cases (Chlamydia)	Rate (Chlamydia)	Cases (Gonorrhea)	Rate (Gonorrhea)
Aitkin	11	68	0	-
Anoka	814	221	194	53
Becker	56	158	6	17
Beltrami	108	231	8	17
Benton	89	215	14	34
Big Stone	5	97	2	-
Blue Earth	157	225	15	22
Brown	21	82	2	-
Carlton	51	139	8	22

County	Cases (Chlamydia)	Rate (Chlamydia)	Cases (Gonorrhea)	Rate (Gonorrhea)
Carver	131	119	20	18
Cass	45	144	4	-
Chippewa	18	147	0	-
Chisago	70	121	6	10
Clay	191	290	44	67
Clearwater	9	104	0	-
Cook	3	-	1	-
Cottonwood	20	176	0	-
Crow Wing	76	112	14	21
Dakota	914	206	237	53
Dodge	40	191	2	-
Douglas	31	78	3	-
Faribault	10	72	3	-
Fillmore	18	84	3	-
Freeborn	54	176	3	-
Goodhue	52	108	13	27
Grant	5	81	1	-
Hennepin	4,580	363	2449	194
Houston	18	96	4	-
Hubbard	22	100	0	-
Isanti	55	129	10	23
Itasca	57	126	1	-
Jackson	9	91	1	-
Kanabec	31	188	2	-
Kandiyohi	78	178	7	16
Kittson	6	148	0	-
Koochiching	9	76	1	-
Lac qui Parle	7	105	0	-
Lake	7	64	1	-
Lake of the Woods	3	-	0	-
Le Sueur	28	96	5	17
Lincoln	5	90	0	-

County	Cases (Chlamydia)	Rate (Chlamydia)	Cases (Gonorrhea)	Rate (Gonorrhea)
Lyon	38	150	0	-
McLeod	44	120	5	14
Mahnomen	17	319	1	-
Marshall	5	56	0	-
Martin	19	97	3	-
Meeker	15	64	3	-
Mille Lacs	25	92	4	-
Morrison	32	93	6	18
Mower	121	301	9	22
Murray	8	99	6	74
Nicollet	34	99	0	-
Nobles	73	333	3	-
Norman	8	125	2	-
Olmsted	312	190	49	30
Otter Tail	50	83	7	12
Pennington	15	108	2	-
Pine	30	102	3	-
Pipestone	10	107	0	-
Polk	46	150	10	33
Pope	12	105	1	-
Ramsey	1,967	367	816	152
Red Lake	4	-	0	-
Redwood	9	59	1	-
Renville	10	69	0	-
Rice	91	134	21	31
Rock	6	63	2	-
Roseau	11	72	4	-
St. Louis	330	165	99	50
Scott	288	186	77	50
Sherburne	176	175	27	27
Sibley	19	127	0	-
Stearns	325	203	43	27

County	Cases (Chlamydia)	Rate (Chlamydia)	Cases (Gonorrhea)	Rate (Gonorrhea)
Steele	62	166	6	16
Stevens	14	145	3	-
Swift	17	174	2	-
Todd	33	129	2	-
Traverse	2	-	0	-
Wabasha	18	83	4	-
Wadena	15	105	2	-
Waseca	19	101	3	-
Washington	438	159	87	32
Watonwan	12	108	3	-
Wilkin	6	94	2	-
Winona	77	156	4	-
Wright	212	143	39	26
Yellow Medicine	9	95	0	-

[l]Rates not calculated for counties with fewer than five cases.

Note: County syphilis data is available at Newly Diagnosed Syphilis Cases by County (https://www.health.state.mn.us/diseases/stds/stats/county.html).

Minnesota Department of Health
651-201-5414
www.health.state.mn.us/std

10/28/2026

To obtain this information in a different format, call: 651-201-5414.